

## Sayana Press Pilot Introduction Project

### Post-training Evaluation

Name: \_\_\_\_\_ Date: \_\_\_\_\_

Region: \_\_\_\_\_ District: \_\_\_\_\_

Question	Score
For questions 1-6, tick all responses that are correct. Some questions have multiple correct answers. If you do not know the correct response, leave the question blank.	
1. Sayana Press prevents pregnancy in women for a period of: <input type="checkbox"/> 6 months <input type="checkbox"/> 3 months <input type="checkbox"/> 1 month only	
2. Which of the following are common side effects of Sayana Press? <input type="checkbox"/> Headache <input type="checkbox"/> Weight changes <input type="checkbox"/> Mood changes and changes in sex drive <input type="checkbox"/> Infertility if used for a long time <input type="checkbox"/> Infrequent or no menstrual bleeding <input type="checkbox"/> Birth defects or miscarriage if the woman is pregnant	
3. Who should not use Sayana Press? <input type="checkbox"/> Women who do not yet have any children <input type="checkbox"/> Women who smoke cigarettes <input type="checkbox"/> Women who are breastfeeding <input type="checkbox"/> Women with high blood pressure <input type="checkbox"/> Women with a history of heart attacks or strokes <input type="checkbox"/> Women with migraine headaches	
4. What are the injection sites for Sayana Press? <input type="checkbox"/> Abdomen <input type="checkbox"/> Back of the arm <input type="checkbox"/> Buttocks <input type="checkbox"/> Front of the thigh <input type="checkbox"/> Deltoid arm muscle	
5. After giving a Sayana Press injection: <input type="checkbox"/> Recap the needle before disposing in the sharps container <input type="checkbox"/> Do not recap the needle and immediately dispose in the sharps container	

6. When is it okay to re-inject a client with Sayana Press without ruling out pregnancy?	
<input type="checkbox"/> When your client arrives for reinjection 2 weeks early <input type="checkbox"/> When your client arrives for reinjection 2 weeks late <input type="checkbox"/> When your client arrives for reinjection 4 weeks late <input type="checkbox"/> When your client arrives for reinjection 6 weeks late	
For questions 7-10, write the correct response in the blank.	
7. Your client is more than 4 weeks late for her reinjection but she is not pregnant. You administer Sayana Press and advise your client to use a back-up method for _____ (#) days in order to avoid becoming pregnant.	
8. A woman using Sayana Press should plan to return for reinjection after how many weeks? _____	
9. The most common side effect that leads women to stop using the injectable is: _____	
For questions 11-16, circle the correct response, either True or False	
10. True / False      The condom is the only method that prevents both pregnancy and HIV.	
11. True / False      Depo IM and Sayana Press have the same active drug, and are equally effective at preventing pregnancy.	
12. True / False      It is acceptable for a client who received Depo IM at her last injection to get Sayana Press for her next injection.	
13. True / False      After stopping Sayana Press, it takes some time before fertility returns, but most women will be able to become pregnant about 10-12 months after their last injection.	
14. True / False      If the provider sees any liquid medication remaining in the reservoir, the injection will not be effective.	
15. True / False      It is safe for women who are breastfeeding to use Sayana Press when their baby is at least 6 weeks old.	
16. True / False      A client only needs counseling before their first injection with Sayana Press.	
	Total Score
	Total possible <b>25</b>
	Percent