Sayana Press Introduction Project

Post-training Evaluation – ANSWER KEY

Name:	Date:
Region:	District:

Question	Score	
For questions 1-6, tick all responses that are correct. Some questions have multiple		
correct answers. If you do not know the correct response, leave the question blank		
(The correct answers for each question are bolded .)		
1. Sayana Press prevents pregnancy in women for a period of:		
□ 6 months	1	
☐ 3 months (1 point)		
☐ 1 month only		
2. Which of the following are common side effects of Sayana Press?		
☐ Headache (1 point)		
☐ Weight changes (1 point)		
☐ Mood changes and changes in sex drive (1 point)	4	
☐ Infertility if used for a long time		
☐ Infrequent or no menstrual bleeding (1 point)		
☐ Birth defects or miscarriage if the woman is pregnant		
(total of 4 points possible)		
3. Who should not use Sayana Press?		
☐ Women who do not yet have any children		
☐ Women who smoke cigarettes		
☐ Women who are breastfeeding	3	
☐ Women with high blood pressure (1 point)		
☐ Women with a history of heart attacks or strokes (1 point)		
☐ Women with migraine headaches (1 point)		
(total of 3 points possible)		
4. What are the injection sites for Sayana Press?		
☐ Abdomen (1 point)		
☐ Back of the arm (1 point)		
□ Buttocks	3	
☐ Front of the thigh (1 point)		
☐ Deltoid arm muscle		
(total of 3 points possible)		
5. After giving a Sayana Press injection:		
 Recap the needle before disposing in the sharps container 		
 Do not recap the needle and immediately dispose in the sharps container 	1	
(1 point)		
6. When is it okay to re-inject a client with Sayana Press without ruling out		
pregnancy?	3	
☐ When your client arrives for reinjection 2 weeks early (1 point)		
☐ When your client arrives for reinjection 2 weeks late (1 point)		
☐ When your client arrives for reinjection 4 weeks late (1 point)		
☐ When your client arrives for reinjection 6 weeks late		
(total of 3 points possible)		

For questions 7-10), write the correct response in the blank.		
 Your client is more than 4 weeks late for her reinjection but she is not pregnant. You administer Sayana Press and advise your client to use a back-up method for (#) days in order to avoid becoming pregnant. Correct Response: 7 (1 point) 		1	
many weeks?	g Sayana Press should plan to return for reinjense: 13 weeks (1 point)	ection after how	1
 The most common side effect that leads women to stop using the injectable is: Correct Response: Menstrual bleeding changes or stops (1 point) 		1	
For questions 11-1	.6 circle the correct response, either True or Fa	alse.	
10. True / False	True / False The condom is the only method that prevents both pregnancy and HIV. (1 point)		1
11. True / False	lse Depo IM and Sayana Press have the same active drug, and are equally effective at preventing pregnancy. (1 point)		1
12. True / False	It is acceptable for a client who received Depo IM at her last injection to get Sayana Press for her next injection. (1 point)		1
13. True / False	rue / False After stopping Sayana Press, it takes some time before fertility returns, but most women will be able to become pregnant about 10-12 months after their last injection. (1 point)		1
14. True / False	4. True / False If the provider sees any liquid medication remaining in the reservoir, the injection will not be effective. (1 point) It is okay if a small amount of liquid remains in the reservoir.		1
15. True / False	5. True / False It is safe for women who are breastfeeding to use Sayana Press when their baby is at least 6 weeks old. (1 point)		1
16. True / False	6. True / False A client only needs counseling before their first injection with Sayana Press. (1 point) Clients should receive counseling at every injection.		1
A recommended passing score is 80% or better, or a minimum of 20 points correct out of 25 points possible. Total Score Total possible		25	
		·	25
		Percent	100%

Directions to the trainer:

Write the number of points scored for each question in the score column to the right of the question. Note that several questions are worth more than one point and that the trainee can get partial credit for ticking some of the correct responses. Write the total score for the post-training evaluation on the total score line and use the total possible score of 25 to calculate the percent correct. A recommended passing score is 80% (20 points) or better.